

August 8, 2025

Name of listed company: Chugai Pharmaceutical Co., Ltd.
Code number: 4519 (Prime Market of Tokyo Stock Exchange)
Head office: 1-1, Nihonbashi-Muromachi 2-Chome, Chuo-ku, Tokyo
President & CEO: Osamu Okuda
Inquiries to: Kae Miyata
Head of Corporate Communications Dept.
Tel: +81-(0)3-3273-0554

**Eli Lilly's Announcement Regarding Oral GLP-1 Orforglipron
(Statistically Significant Weight Reduction and a Safety Profile
Consistent with Injectable GLP-1 Medicines in Phase III ATTAIN-1
in Adults with Obesity)**

TOKYO, August 8, 2025 – [Chugai Pharmaceutical Co., Ltd.](#) (TOKYO: 4519) announced that [Eli Lilly and Company](#) (hereafter “Eli Lilly”, NYSE: LLY) issued a press release on August 7, 2025 (local time) regarding the positive topline Phase 3 results from ATTAIN-1, evaluating the efficacy and safety of orforglipron compared to placebo in adults with obesity, or overweight with a weight-related medical problem and without diabetes.

In the first Phase 3 trial of the ATTAIN program, ATTAIN-1 trial (n=3,127), orforglipron, taken once per day without food and water restrictions, achieved the primary and all key secondary endpoints compared to placebo. Participants taking the highest dose of orforglipron lost an average of 27.3 lbs (12.4kg) (12.4%) at 72 weeks using the efficacy estimand.* In a key secondary endpoint, 59.6% of participants taking the highest dose of orforglipron lost at least 10% of their body weight, while 39.6% lost at least 15% of their body weight.

The overall safety profile of orforglipron in ATTAIN-1 was consistent with the established GLP-1 receptor agonist class. The most commonly reported adverse events were gastrointestinal-related (nausea, constipation, diarrhea, vomiting, dyspepsia) and generally mild to moderate in severity. Treatment discontinuation rates due to adverse events were 5.1% (6 mg), 7.7% (12 mg) and 10.3% (36 mg) for orforglipron vs. 2.6% with placebo. The overall treatment discontinuation rates were 21.9% (6 mg), 22.5% (12 mg) and 24.4% (36 mg) for orforglipron vs. 29.9% with placebo. No hepatic safety signal was observed.

The detailed ATTAIN-1 results will be presented next month at the European Association for the Study of Diabetes (EASD) Annual Meeting 2025 and published in a peer-reviewed journal. More results from the ATTAIN Phase 3 clinical trial program will be shared later this year, along with findings from the ACHIEVE Phase 3 clinical trial program evaluating orforglipron for adults with type 2 diabetes. Eli Lilly expects to submit orforglipron for weight management to global regulatory agencies by the end of this year.

For the full text of Eli Lilly's press release, please refer to the Latest News via the following link:
Lilly's oral GLP-1, orforglipron, delivers weight loss of up to an average of 27.3 lbs in first of
two pivotal Phase 3 trials in adults with obesity

<https://investor.lilly.com/>

This announcement is not expected to have an impact on Chugai's consolidated financial forecast
for the fiscal year ending December 2025, which was announced on January 30, 2025.

* The efficacy estimand represents efficacy had all randomized participants remained on
study intervention (with possible dose interruptions and modifications) for 72 weeks
without initiating prohibited weight management treatments.

###